

Range-Separated Random Phase Approximations

Bastien Mussard^{1,2}

J. Toulouse¹, P. Reinhardt¹, O. Franck¹, E. Luppi¹,
D. Rocca³, J. G. Ángyán³

¹ Laboratoire de Chimie Théorique, Université Pierre et Marie Curie

² Institut du Calcul et de la Simulation, Université Pierre et Marie Curie

³ Cristallographie Résonance Magnétique et Modélisation, Université de Lorraine

bastien.mussard@upmc.fr

www.lct.jussieu.fr/pagesperso/mussard/

Adiabatic connection formula for E_c

$$E_c = \frac{1}{2} \int_0^1 d\alpha \operatorname{Tr}(\hat{W}_{ee} \cdot \mathbb{P}_{c,\alpha}) = \int_0^1 d\alpha (\langle \Psi_\alpha | \hat{W}_{ee} | \Psi_\alpha \rangle - \langle \Phi_0 | \hat{W}_{ee} | \Phi_0 \rangle)$$

Fluctuation-dissipation theorem for $P_{c,\alpha}$

$$\mathbb{P}_{c,\alpha} = \int_{-\infty}^{\infty} \frac{-d\omega}{2\pi} [\mathbb{\Pi}_\alpha(i\omega) - \mathbb{\Pi}_0(i\omega)]$$

Linear response equations for $\mathbb{\Pi}_\alpha(i\omega)$

$$\mathbb{\Pi}_\alpha(i\omega)^{-1} = \mathbb{\Pi}_0(i\omega)^{-1} - \mathbb{f}_{\text{Hxc},\alpha}(i\omega)$$

RPA correlation energy $\mathbb{f}_{\text{xc},\alpha} = 0$ or $\mathbb{f}_{c,\alpha} = 0$

$$E_c = \frac{1}{2} \int_0^1 d\alpha \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \operatorname{Tr}(\hat{W}_{ee} \cdot [\mathbb{\Pi}_\alpha^{\text{RPA}}(i\omega) - \mathbb{\Pi}_0(i\omega)])$$

where $\mathbb{\Pi}_\alpha^{\text{RPA}}$ is obtained from the eigenvectors of $\begin{pmatrix} \mathbf{A}_\alpha & \mathbf{B}_\alpha \\ -\mathbf{B}_\alpha & -\mathbf{A}_\alpha \end{pmatrix}$

RPA : different formalisms

$$E_c = \frac{1}{2} \int_0^1 d\alpha \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \text{Tr} (W_{ee} \cdot [\Pi_{\alpha}^{\text{RPA}}(i\omega) - \Pi_0(i\omega)]) \quad \begin{aligned} (\mathbf{B})_{ia,jb} &= \langle ab|ij\rangle \\ (\mathbf{A})_{ia,jb} &= \Delta\epsilon_{ia,jb} + \langle ib|aj\rangle \end{aligned}$$

- ▶ Analytical integration on the **frequency**

density matrix formulation

$$E_c = \frac{1}{2} \int_0^1 d\alpha \text{Tr} (W_{ee} \cdot P_{c,\alpha}^{\text{RPA}})$$

- ▶ **Both** analytical integration

plasmon formulation

$$E_c = \frac{1}{2} \text{tr} (\Omega^{\text{RPA}} - \Omega^{\text{TDA}})$$

- ▶ which is **equivalent** to the

rCCD formulation

$$E_c = \frac{1}{2} \text{tr} (\mathbf{B} \cdot \mathbf{T})$$

where \mathbf{T} are solutions of the **Riccati** equations

$$\mathbf{B} + \mathbf{AT} + \mathbf{TA} + \mathbf{TBT} = \mathbf{0}$$

[Scuseria,Henderson,Sorensen JCP **129** 231101 (2008)]

- ▶ Analytical integration on the **coupling constant**

dielectric matrix formulation

$$E_c = \frac{1}{2} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \text{tr} (\log [\mathbf{1} - \Pi_0 \mathbf{B}] + \Pi_0 \mathbf{B})$$

[Jansen,BM,Rocca,Ángyán (prep.)]

RPA : different flavors

$$E_c = \frac{1}{2} \text{tr}(\mathbf{B} \cdot \mathbf{T})$$

$$\text{where } \mathbf{B} + \mathbf{A}\mathbf{T} + \mathbf{T}\mathbf{A} + \mathbf{T}\mathbf{B}\mathbf{T} = \mathbf{0}$$

$$(\mathbf{B})_{ia,jb} = \langle ab|ij \rangle$$

$$(\mathbf{A})_{ia,jb} = \Delta\epsilon_{ia,jb} + \langle ib|aj \rangle$$

The Riccati equations, in this context, are derived from $\begin{pmatrix} \mathbf{A} & \mathbf{B} \\ -\mathbf{B} & -\mathbf{A} \end{pmatrix}$

which depend on a kernel \mathbb{f} that **does not or does include exchange**

► **drCCD** variant $\longrightarrow \mathbb{f}_{xc} = 0 \longrightarrow$

$$\begin{aligned} (\mathbf{B}^I)_{ia,jb} &= \langle ab|ij \rangle \\ (\mathbf{A}^I)_{ia,jb} &= \Delta\epsilon_{ia,jb} + \langle ib|aj \rangle \end{aligned}$$

$$E_c^{\text{drRPA-I}} = \frac{1}{2} \text{tr}(\mathbf{B}^I \cdot \mathbf{T}^{\text{drRPA}})$$

$$\text{where } \mathbf{B}^I + \mathbf{A}^I\mathbf{T} + \mathbf{T}\mathbf{A}^I + \mathbf{T}\mathbf{B}^I\mathbf{T} = \mathbf{0}$$

► **rCCD** variant $\longrightarrow \mathbb{f}_c = 0 \longrightarrow$

$$\begin{aligned} (\mathbf{B}^{II})_{ia,jb} &= \langle ab||ij \rangle \\ (\mathbf{A}^{II})_{ia,jb} &= \Delta\epsilon_{ia,jb} + \langle ib||aj \rangle \end{aligned}$$

$$E_c^{\text{RPAx-II}} = \frac{1}{4} \text{tr}(\mathbf{B}^{II} \cdot \mathbf{T}^{\text{RPAx}})$$

$$\text{where } \mathbf{B}^{II} + \mathbf{A}^{II}\mathbf{T} + \mathbf{T}\mathbf{A}^{II} + \mathbf{T}\mathbf{B}^{II}\mathbf{T} = \mathbf{0}$$

► Additional flavors

$$E_c^{\text{SOSEX}} = \frac{1}{2} \text{tr}(\mathbf{B}^{II} \cdot \mathbf{T}^{\text{drRPA}})$$

$$E_c^{\text{SO2}} = \frac{1}{2} \text{tr}(\mathbf{B}^I \cdot \mathbf{T}^{\text{RPAx}})$$



Note that comparable additional flavors also exist in the **density matrix** formulation (they have been explored in the literature) and in the **dielectric matrix** formulation.

DFT calculation followed by an **RPA calculation using KS orbitals and energies**

$$E^{\text{DFT}} = \min_{\Phi} \left\{ \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} | \Phi \rangle + E_{\text{Hxc}}[n_{\Phi}] \right\}$$

$$E = \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} | \Phi \rangle + E_{\text{Hxc}}^{\text{HF}}[\Phi] + E_{\text{c}}^{\text{RPA}}$$

- ▶ DFT (LDA/GGA/...) do not describe well the **long-range dispersion forces**, while RPA does
- ▶ Still problems with : **short-range correlation energies** (far too negative)
strong dependence on basis size
simple van der Waals dimers



Range-Separated Hybrid + RPA



- ▶ The idea : split the e-e interaction into a **short-** and a **long-range** parts

Hybrid **DFT** with exact HF exchange and RPA correlation, both at **long-range** :

$$E = \min_{\Phi} \left\{ \langle \Phi | \hat{T} + \hat{V}_{ne} + \hat{W}_{ee}^{lr} | \Phi \rangle + E_{Hxc}^{sr}[n_{\Phi}] \right\} + E_c^{lr,RPA}$$

[Stoll,Savin DF.meth.phys. 177 (1985)][Savin IJQC 22 59 (1988)][Savin, Rec.dev. 327 (1996)]

- ▶ range separation greatly **improves RPA** (see **srPBE+RPAx-SO2**)
- ▶ basis dependence is reduced



>srPBE, aug-cc-pV6Z, $\mu = 0.5$ [Toulouse,Zhu,Savin,Jansen,Ángyán 135 084119 (2011)]

Range-Separated Hybrid + RPA


- ▶ The idea : split the e-e interaction into a **short-** and a **long-range** parts

Hybrid **DFT** with exact HF exchange and RPA correlation, both at **long-range** :

$$E = \min_{\Phi} \left\{ \langle \Phi | \hat{T} + \hat{V}_{ne} + \hat{W}_{ee}^{lr} | \Phi \rangle + E_{Hxc}^{sr}[n_{\Phi}] \right\} + E_c^{lr,RPA}$$

[Stoll,Savin DF.meth.phys. 177 (1985)][Savin IJQC 22 59 (1988)][Savin, Rec.dev. 327 (1996)]

- ▶ range separation greatly **improves RPA** (see **srPBE+RPAx-SO2**)
- ▶ basis dependence is reduced
- ▶ for ex. : performance on the S22 dataset is very good for **srPBE+RPAx-SO2**






>srPBE, aug-cc-pVDZ, $\mu = 0.5$ [Toulouse,Zhu,Savin,Jansen,Ángyán 135 084119 (2011)]

Wavefunction basis convergence

[Gori-Giorgi,Savin PRA 73 032506 (2006)]

both **partial wave** and **principal number** expansions
lead to **exponential convergence** of the wavefunction



Extrapolation formula for total energies

We propose a **three-point extrapolation scheme** for total RSH+correlation energies ($E_X = E_\infty + B \exp(-\beta X)$)

$$E_\infty = E_{XYZ} = \frac{E_Y^2 - E_X E_Z}{2E_Y - E_X - E_Z}$$

>errors (mHartree) wrt. cc-pV6Z

	He	Ne	N ₂	H ₂ O
ΔE_D	8.488	74.523	51.581	55.850
ΔE_T	0.781	20.337	13.406	14.736
ΔE_Q	0.245	5.763	4.090	4.499
ΔE_5	0.078	0.751	0.810	0.726
ΔE_{DTQ}	0.205	0.401	1.083	1.105
ΔE_{TQ5}	0.002	-1.876	-0.972	-1.475


Implemented most of the variants with a nospinflip/spinflip block structure.

Normal distributions of errors (kcal/mol) of calculations on **AE49**


- mean errors of post-RSH calculation are better
- post-RSH calculations dist. of errors have **sharper distributions**

Similar results were obtained on the **DBH24/08** dataset


We argue that **RPA_x-SO2** is a good method for a wide range of applications




HF+MP2





RSH+MP2



HF+dRPA-I



RSH+dRPA-I

HF+RPA_x-SO2RSH+RPA_x-SO2

>cc-pVQZ (post-RSH : srPBE, $\mu = 0.5$)

Fractional Occupation Numbers calculations

- Study of systems with a **non-integer charge**

$$E^{N+\delta} = \min_{\hat{f} \rightarrow N+\delta} \text{Tr} \left[\hat{f} \left(\hat{T} + \hat{V}_{\text{ext}} + \hat{V}_{\text{ee}} \right) \right]$$

$$\hat{f}^{N+\delta} = (1 - \delta)\hat{f}^N + (\delta)\hat{f}^{N+1}$$

$$E^{N+\delta} = (1 - \delta)E^N + (\delta)E^{N+1}$$




$$n^{N+\delta} = (1 - \delta)n^N + (\delta)n^{N+1}$$

E is **piecewise linear** wrt N and has a **derivative discontinuity** at integer N

$$E^{N+\delta} = \min_{\hat{f} \rightarrow N+\delta} \text{Tr} \left[\hat{f} \left(\hat{T} + \hat{V}_{\text{ext}} \right) + E_{\text{Hxc}}[n_{\hat{f}}] \right]$$


$$\hat{f}_s^{N+\delta} = (1 - \delta)\hat{f}_s^{N,\delta} + (\delta)\hat{f}_n^{N+1,\delta}$$

$$n^{N+\delta}(\mathbf{r}) = \sum f_i |\phi_i^{N+\delta}(\mathbf{r})|^2$$




Fractional Occupation Numbers calculations

- ▶ Study of systems with a **non-integer spin**




- ▶ Systems with fractional occupation of degenerate spin states should have the same energy as the integer-spin states





- ▶ **Constancy condition**

$$E \left[\sum c_i n_i \right] = E[n] = E(N)$$




Fractional Occupation Numbers calculations [early results]




- PBE+RPA - - - ● - - -
- srPBE+lrRPA - - - ● - - -
- PBE+SO2 - - - ■ - - -
- srPBE+lrSO2 - - - ■ - - -
- PBE+SOSEX - - - ◆ - - -
- srPBE+lrSOSEX - - - ◆ - - -

> deviation from linear extrapolation for H
with δ electron, cc-pVDZ




(same thing for He)











- PBE+RPA ● - - -
- srPBE+lrRPA ● —
- PBE+SO₂ ■ - - -
- srPBE+lrSO₂ ■ —
- PBE+SOSEX ◆ - - -
- srPBE+lrSOSEX ◆ —

> deviation from linear extrapolation for He
with $1 + \delta$ electrons, cc-pVDZ




Fractional Occupation Numbers calculations [early results]



PBE+RPA 
srPBE+lrRPA 
PBE+SO₂ 
srPBE+lrSO₂ 
PBE+SOSEX 
srPBE+lrSOSEX 

δ energy for H using cc-pVDZ
 with $(0.5 + \delta)$ α -electron and $(0.5 - \delta)$ β -electron



Conclusions (and future work)

- ▶ All (and more) is implemented in the **Quantum Chemistry Package MOLPRO**
Forces for (RSH+)RPA energies are available
and so are **geometry optimizations** at the (RSH+)RPA levels
[BM,Szalay,Ángyán JCTC **10** 1968 (2014)]
- ▶ Using the fractional occupations implementations
to get IP and AE with double-hybrids
- ▶ Instabilities in the RPA problem
- ▶ at the Université Pierre et Marie Curie
Odile Franck
Eleanora Luppi
Peter Reinhardt
Julien Toulouse
and, at the Université de Lorraine
János Ángyán
Dario Rocca